




TRP14 Monoclonal Antibody

| Catalog No | BYmab-02855 |
|---------------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | TXNDC17 |
| Protein Name | Thioredoxin domain-containing protein 17 |
| Immunogen | Synthesized peptide derived from TRP14 . at AA range: 20-100 |
| Specificity | TRP14 Monoclonal Antibody detects endogenous levels of TRP14 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | TXNDC17; TXNL5; Thioredoxin domain-containing protein 17; 14 kDa thioredoxin-related protein; TRP14; Protein 42-9-9; Thioredoxin-like protein 5 |
| Observed Band | 13kD |
| | |
| Cell Pathway | Cytoplasm . |
| Cell Pathway Tissue Specificity | Cytoplasm . Ubiquitously expressed in cell lines. |
| | Ubiquitously expressed in cell lines. function:Disulfide reductase. May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and |
| Tissue Specificity | Ubiquitously expressed in cell lines. function:Disulfide reductase. May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and NF-kappa-B activation. Has peroxidase activity and may contribute to the elimination of cellular hydrogen peroxide.,PTM:The oxidized protein is reduced by TRXR1.,similarity:Belongs to the thioredoxin family.,similarity:Contains 1 thioredoxin domain.,subunit:Interacts with TRXR1 and DYNLL1/DNCL1.,tissue |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and NF-kappa-B activation. Has peroxidase activity and may contribute to the elimination of cellular hydrogen peroxide.,PTM:The oxidized protein is reduced by TRXR1.,similarity:Belongs to the thioredoxin family.,similarity:Contains 1 thioredoxin domain.,subunit:Interacts with TRXR1 and DYNLL1/DNCL1.,tissue specificity:Ubiquitously expressed in cell lines.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TRP14 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd