



HDC Monoclonal Antibody

Catalog No	BYmab-02862
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HDC
Protein Name	Histidine decarboxylase
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human HDC. AA range:201-250
Specificity	HDC Monoclonal Antibody detects endogenous levels of HDC protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HDC; Histidine decarboxylase; HDC
Observed Band	72kD
Cell Pathway	cytosol,
Tissue Specificity	Leukemia,
Function	catalytic activity:L-histidine = histamine + CO(2).,cofactor:Pyridoxal phosphate.,pathway:Amine and polyamine biosynthesis; histamine biosynthesis; histamine from L-histidine: step 1/1.,similarity:Belongs to the group II decarboxylase family.,subunit:Homodimer.,
Background	This gene encodes a member of the group II decarboxylase family and forms a homodimer that converts L-histidine to histamine in a pyridoxal phosphate dependent manner. Histamine regulates several physiologic processes, including neurotransmission, gastric acid secretion,inflammation, and smooth muscle tone.[provided by RefSeq, Aug 2010],

Nanjing BYabscience technology Co.,Ltd




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using HDC Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658