



## GCP5 Monoclonal Antibody

Catalog No         BYmab-03138           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         TUBGCP5           Protein Name         Gamma-tubulin complex component 5           Immunogen         The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range;741-790           Specificity         GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5           Observed Band         118kD           Cell Pathway         Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .           Tissue Specificity         Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Function         dunction:Gamma-tubulin complex		
Reactivity Human;Mouse;Rat  Applications WB  Gene Name TUBGCP5  Protein Name Gamma-tubulin complex component 5  Immunogen The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range:741-790  Specificity GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background	Catalog No	BYmab-03138
Applications  Gene Name  TUBGCP5  Protein Name  Gamma-tubulin complex component 5  Immunogen  The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range:741-780  Specificity  GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse,IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band  118kD  Cell Pathway  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity-Belongs to the GCP family, subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity: Belongs to the GCP family, subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity: Belongs to the GCP family, subunit: Gamma-tubulin complex is scomposed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in heart and highest levels in heart and skeletal muscle and moderate levels in heart and skeletal muscle	Isotype	IgG
Gene Name         TUBGCP5           Protein Name         Gamma-tubulin complex component 5           Immunogen         The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range:741-790           Specificity         GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5           Observed Band         118kD           Cell Pathway         Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .           Tissue Specificity         Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Function         function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity. Belongs to the GCP family, subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, Issues specificity. Widely expressed, with highest levels in heart and GCP6, Issues specificity: W	Reactivity	Human;Mouse;Rat
Protein Name         Gamma-tubulin complex component 5           Immunogen         The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range:741-790           Specificity         GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5           Observed Band         118kD           Cell Pathway         Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .           Tissue Specificity         Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Function         function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from human TUBGCP5. AA range:741-790           Specificity         GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5           Observed Band         118kD           Cell Pathway         Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.           Tissue Specificity         Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Function         function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Background         function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, si	Gene Name	TUBGCP5
TUBGCP5. AA range:741-790  Specificity GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome. similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity. Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.	Protein Name	Gamma-tubulin complex component 5
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family, .subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family, .subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family, .subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, .similarity:Belongs to the GCP family in the gCP3 GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.	Immunogen	
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5           Observed Band         118kD           Cell Pathway         Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .           Tissue Specificity         Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.           Function         function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and GCP6, tissue specificity:Widely expressed by the highest levels in heart and	Specificity	GCP5 Monoclonal Antibody detects endogenous levels of GCP5 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5 Observed Band 118kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome. similarity:Belongs to the GCP family., subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family., subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family subunit:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP family subunit:Gamma-tubulin complex	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Widely expressed, with highest levels in heart and GCP6, tissue specificity:Widely expressed, with highest levels in heart and	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band 118kD  Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .  Tissue Specificity Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Dilution	WB 1:500-2000
Synonyms  TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band  118kD  Cell Pathway  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Widely expressed, with highest levels in heart and GCP6, tissue specificity: Widely expressed, with highest levels in heart and	Concentration	1 mg/ml
Synonyms  TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5  Observed Band  118kD  Cell Pathway  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background  function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Widely expressed, with highest levels in heart and	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and		
Tissue Specificity  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  Function  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Synonyms	TUBGCP5; GCP5; KIAA1899; Gamma-tubulin complex component 5; GCP-5
Function  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	•	
centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.,  Background  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Observed Band	118kD
centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and	Observed Band Cell Pathway	118kD  Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate
	Observed Band Cell Pathway Tissue Specificity	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.  Widely expressed, with highest levels in heart and skeletal muscle and moderate levels in brain.  function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Widely expressed, with highest levels in heart and

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using GCP5 Monoclonal Antibody