



MYBPC1 Monoclonal Antibody

| Catalog No | BYmab-03154 |
|--------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | MYBPC1 |
| Protein Name | Myosin-binding protein C slow-type |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MYBPC1. AA range:218-267 |
| Specificity | MYBPC1 Monoclonal Antibody detects endogenous levels of MYBPC1 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,lgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | MYBPC1; MYBPCS; Myosin-binding protein C; slow-type; Slow MyBP-C; C-protein, skeletal muscle slow isoform |
| Observed Band | 120kD |
| Cell Pathway | cytosol,myofibril,myosin filament, |
| Tissue Specificity | Brain,Fetal skeletal muscle,Hippocampus,Liver,Muscle pool- 2 tissue |
| Function | function:Thick filament-associated protein located in the crossbridge region of vertebrate striated muscle a bands. In vitro it binds MHC, F-actin and native thin filaments, and modifies the activity of actin-activated myosin ATPase. It may modulate muscle contraction or may play a more structural role.,similarity:Belongs to the immunoglobulin superfamily. MyBP family.,similarity:Contains 3 fibronectin type-III domains.,similarity:Contains 7 Ig-like C2-type (immunoglobulin-like) domains., |
| Background | This gene encodes a member of the myosin-binding protein C family. Myosin-binding protein C family members are myosin-associated proteins found in the cross-bridge-bearing zone (C region) of A bands in striated muscle. The |
| | Nanjing Byabscience technology Co. Ltd |

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| encoded protein is the slow skeletal muscle isoform of myosin-binding protein and plays an important role in muscle contraction by recruiting muscle-type creatine kinase to myosin filaments. Mutations in this gene are associated with distal arthrogryposis type I. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011], | 1 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|
| A () () () () () | |


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using MYBPC1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd