



Tubulin γ Monoclonal Antibody

Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytocytoskeleton, spindle. Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly, mass spectrometry: PubMed:11840567, similarity. B to the tubulin family, subunit: Interacts with GCP2 and GCP3. Interacts with GCP3 and GCP3. Interacts wit		
Reactivity Human;Mouse;Rat Applications WB Gene Name TUBG1 Protein Name Tubulin gamma-1 chain Immunogen The antiserum was produced against synthesized peptide derived from human Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide and the protein. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the	Catalog No	BYmab-03207
Applications WB Gene Name TUBG1 Protein Name Tubulin gamma-1 chain Immunogen The antiserum was produced against synthesized peptide derived from hu Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide and contains and conta	Isotype	IgG
Gene Name TUBG1 Protein Name Tubulin gamma-1 chain Immunogen The antiserum was produced against synthesized peptide derived from hu Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function:Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule assembly, mass spectrometry: PubMed:11840567, similarity.B to the tubulin family, subunit:Interacts with GCP2 and GCP3. Interacts with to the tubulin superfamily. The encoded proterior for the tubulin family	Reactivity	Human;Mouse;Rat
Protein Name Tubulin gamma-1 chain Immunogen The antiserum was produced against synthesized peptide derived from hu Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulic complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer,Skin, Function function:Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly, mass spectrometry: PubMed:11840567, similarity.B to the tubulin family.subulunit interacts with GCP2 and GCP3. Interacts w	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from hu Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulic complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly, mass spectrometry: PubMed:11840567, similarity. B to the tubulin family, subunit-interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded preferred to as the gamma-tubulin ring complex. The protein mediates microtered to as the gamma-tubulin ring complex. The protein mediates micro	Gene Name	TUBG1
Tubulin gamma. AA range:386-435 Specificity Tubulin γ Monoclonal Antibody detects endogenous levels of Tubulin γ protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer,Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly, mass spectrometry: PubMed:11840567, similarity. B to the tubulin family, subunit:Interacts with GCP2 and GCP3. Interacts wit	Protein Name	Tubulin gamma-1 chain
protein. Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules Tissue Specificity Mammary cancer,Skin, Function function:Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule asspartly. But the tubulin family.,subunit:Interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a complex for the centrosome where it binds to microtubules as part of a complex complex complex complex complex complex contents with microtubules as part of a complex co	Immunogen	The antiserum was produced against synthesized peptide derived from human Tubulin gamma. AA range:386-435
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer,Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly, mass spectrometry. PubMed: 11840567, similarity: B to the tubulin family., subunit: Interacts with GCP2 and GCP3. Interacts with GCP2 and GCP3. Interacts with GCP2 and GCP3. Interacts with localizes to the centrosome where it binds to microtubules as part of a correferred to as the gamma-tubulin ring complex. The protein mediates microtuple as a part of a correferred to as the gamma-tubulin ring complex. The protein mediates microtuple.	Specificity	
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly. mass spectrometry: PubMed:11840567, similarity:B to the tubulin family. subunit:Interacts with GCP2 and GCP3. Interacts with Background This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a correferred to as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex. The protein mediates microtubules as the gamma-tubulin ring complex.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity: B to the tubulin family., subunit: Interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a conference to as the gamma-tubulin ring complex. The protein mediates microferred to as the gamma-tubulin ring complex. The protein mediates microferred to as the gamma-tubulin ring complex. The protein mediates microferred to as the gamma-tubulin ring complex. The protein mediates microferred to as the gamma-tubulin ring complex. The protein mediates microferred to as the gamma-tubulin ring complex.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubuling complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytocytoskeleton, spindle. Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer,Skin, Function function:Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly.,mass spectrometry: PubMed:11840567,similarity:B to the tubulin family.,subunit:Interacts with GCP2 and GCP3. Interacts with Background This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a correferred to as the gamma-tubulin ring complex. The protein mediates microtubules microtubules.	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle . Localizes to mitotic spindle microtubules Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly. mass spectrometry: PubMed: 11840567, similarity: B to the tubulin family. subunit: Interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded proceedings of the centrosome where it binds to microtubules as part of a confedered to as the gamma-tubulin ring complex. The protein mediates micro	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band 51kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity: B to the tubulin family., subunit:Interacts with GCP2 and GCP3. Interacts with Background This gene encodes a member of the tubulin superfamily. The encoded profedered to as the gamma-tubulin ring complex. The protein mediates micro	Concentration	1 mg/ml
Synonyms TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1 Observed Band Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytocytoskeleton, spindle. Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, Function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity:B to the tubulin family., subunit: Interacts with GCP2 and GCP3. Interacts with Interacts with GCP2 and GCP3. Interacts with Interacts to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtubules as part of a complex to the centrosome where it binds to microtub	Purity	≥90%
Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytocytoskeleton, spindle . Localizes to mitotic spindle microtubules. Tissue Specificity Mammary cancer, Skin, function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity:B to the tubulin family., subunit:Interacts with GCP2 and GCP3. Interacts with Background This gene encodes a member of the tubulin superfamily. The encoded professional profession of the centrosome where it binds to microtubules as part of a compresented to as the gamma-tubulin ring complex. The protein mediates micro	Storage Stability	-20°C/1 year
Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytocytoskeleton, spindle. Localizes to mitotic spindle microtubules Tissue Specificity Mammary cancer, Skin, function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity: B to the tubulin family., subunit: Interacts with GCP2 and GCP3. Interacts with Chapter of the tubulin superfamily. The encoded process of the centrosome where it binds to microtubules as part of a complex of the protein mediates microtubules.	Synonyms	TUBG1; TUBG; Tubulin gamma-1 chain; Gamma-1-tubulin; Gamma-tubulin complex component 1; GCP-1
Tissue Specificity Mammary cancer, Skin, function function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., mass spectrometry: PubMed:11840567, similarity: B to the tubulin family., subunit: Interacts with GCP2 and GCP3. Interacts with Background This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a comreferred to as the gamma-tubulin ring complex. The protein mediates micro	Observed Band	51kD
function: Tubulin is the major constituent of microtubules. Gamma tubulin is at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly.,mass spectrometry: PubMed:11840567,similarity:B to the tubulin family.,subunit:Interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded process to the centrosome where it binds to microtubules as part of a comreferred to as the gamma-tubulin ring complex. The protein mediates micro	Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle . Localizes to mitotic spindle microtubules
at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly.,mass spectrometry: PubMed:11840567,similarity:B to the tubulin family.,subunit:Interacts with GCP2 and GCP3. Interacts with This gene encodes a member of the tubulin superfamily. The encoded prolocalizes to the centrosome where it binds to microtubules as part of a comreferred to as the gamma-tubulin ring complex. The protein mediates micro	Tissue Specificity	Mammary cancer,Skin,
localizes to the centrosome where it binds to microtubules as part of a com referred to as the gamma-tubulin ring complex. The protein mediates micro	Function	function:Tubulin is the major constituent of microtubules. Gamma tubulin is found at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly.,mass spectrometry: PubMed:11840567,similarity:Belongs to the tubulin family.,subunit:Interacts with GCP2 and GCP3. Interacts with B9D2.,
nucleation and is required for microtubule formation and progression of the	Background	This gene encodes a member of the tubulin superfamily. The encoded protein localizes to the centrosome where it binds to microtubules as part of a complex referred to as the gamma-tubulin ring complex. The protein mediates microtubule nucleation and is required for microtubule formation and progression of the cell


Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	cycle. A pseudogene of this gene is found on chromosome 7. [provided by RefSeq, Jan 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658