



# MB67 Monoclonal Antibody

<b>Catalog No</b>	BYmab-03315
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	NR1I3
<b>Protein Name</b>	Nuclear receptor subfamily 1 group I member 3
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human NR1I3. AA range:71-120
<b>Specificity</b>	MB67 Monoclonal Antibody detects endogenous levels of MB67 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	NR1I3; CAR; Nuclear receptor subfamily 1 group I member 3; Constitutive activator of retinoid response; Constitutive active response; Constitutive androstane receptor; CAR; Orphan nuclear receptor MB67
<b>Observed Band</b>	35kD
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Cytoplasm, cytoskeleton. Recruited to the cytoplasm by DNAJC7. .
<b>Tissue Specificity</b>	Predominantly expressed in liver.
<b>Function</b>	domain:Composed by a short N-terminal domain followed by the DNA binding, hinge, and ligand binding/dimerization domains.,function:Binds and transactivates the retinoic acid response elements that control expression of the retinoic acid receptor beta 2 and alcohol dehydrogenase 3 genes. Transactivates both the phenobarbital responsive element module of the human CYP2B6 gene and the CYP3A4 xenobiotic response element.,induction:By dexamethasone.,similarity:Belongs to the nuclear hormone receptor family.,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding

**Nanjing BYabscience technology Co.,Ltd**



domain.,subunit:Heterodimer of NR1I3 and RXR. Interacts with PSMC4.,tissue specificity:Predominantly expressed in liver.,

#### Background

This gene encodes a member of the nuclear receptor superfamily, and is a key regulator of xenobiotic and endobiotic metabolism. The protein binds to DNA as a monomer or a heterodimer with the retinoid X receptor and regulates the transcription of target genes involved in drug metabolism and bilirubin clearance, such as cytochrome P450 family members. Unlike most nuclear receptors, this transcriptional regulator is constitutively active in the absence of ligand but is regulated by both agonists and inverse agonists. Ligand binding results in translocation of this protein to the nucleus, where it activates or represses target gene transcription. These ligands include bilirubin, a variety of foreign compounds, steroid hormones, and prescription drugs. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Nanjing BYabscience technology Co.,Ltd