



NOR-1 Monoclonal Antibody

Catalog No	BYmab-03321
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NR4A3
Protein Name	Nuclear receptor subfamily 4 group A member 3
Immunogen	The antiserum was produced against synthesized peptide derived from human NR4A3. AA range:387-436
Specificity	NOR-1 Monoclonal Antibody detects endogenous levels of NOR-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NR4A3; CHN; CSMF; MINOR; NOR1; TEC; Nuclear receptor subfamily 4 group A member 3; Mitogen-induced nuclear orphan receptor; Neuron-derived orphan receptor 1; Nuclear hormone receptor NOR-1
Observed Band	65kD
Cell Pathway	Nucleus .
Tissue Specificity	Isoform alpha is highly expressed in skeletal muscle. Isoform beta is highly expressed in skeletal muscle and low expressed in fetal brain and placenta.
Function	disease:A chromosomal aberration involving NR4A3 is a cause of a form of extraskelatal myxoid chondrosarcomas (EMC). Translocation t(9;17)(q22;q11) with TAF2N.,disease:A chromosomal aberration involving NR4A3 is a cause of Ewing sarcoma [MIM:133450]. Translocation t(9;22)(q22-31;q11-12) with EWS.,function:Binds to the B1A response-element.,similarity:Belongs to the nuclear hormone receptor family.,similarity:Belongs to the nuclear hormone receptor family. NR4 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,tissue specificity:High expression of isoform alpha in skeletal muscle. High expression of isoform beta in skeletal muscle and low

Nanjing BYabscience technology Co.,Ltd



expression in fetal brain and placenta.,

Background

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcriptional activator. The protein can efficiently bind the NGFI-B Response Element (NBRE). Three different versions of extraskeletal myxoid chondrosarcomas (EMCs) are the result of reciprocal translocations between this gene and other genes. The translocation breakpoints are associated with Nuclear Receptor Subfamily 4, Group A, Member 3 (on chromosome 9) and either Ewing Sarcome Breakpoint Region 1 (on chromosome 22), RNA Polymerase II, TATA Box-Binding Protein-Associated Factor, 68-KD (on chromosome 17), or Transcription factor 12 (on chromosome 15). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NOR-1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd