



# FA8A (heavy chain 92k, Cleaved-Arg759) mouse mAb

<b>Catalog No</b>	BYmab-03361
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	F8 F8C
<b>Protein Name</b>	FA8A (heavy chain 92k, Cleaved-Arg759)
<b>Immunogen</b>	Synthesized peptide derived from human FA8A (heavy chain 92k, Cleaved-Arg759)
<b>Specificity</b>	This antibody detects endogenous levels of Human,Mouse FA8A (heavy chain 92k, Cleaved-Arg759, protein was cleaved amino acid sequence between 759-760 )
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Coagulation factor VIII (Antihemophilic factor;AHF;Procoagulant component) [Cleaved into: Factor VIIIa heavy chain, 200 kDa isoform; Factor VIIIa heavy chain, 92 kDa isoform; Factor VIII B chain; Factor VIIIa light chain]
<b>Observed Band</b>	92 260kD
<b>Cell Pathway</b>	Secreted, extracellular space.
<b>Tissue Specificity</b>	
<b>Function</b>	acute inflammatory response, defense response, acute-phase response, inflammatory response, cell adhesion, blood coagulation, hemostasis, response to wounding, biological adhesion, wound healing, coagulation, regulation of body fluid levels, oxidation reduction,
<b>Background</b>	disease:Defects in F8 are the cause of hemophilia A (HEMA) [MIM:306700]. HEMA is a common recessive X-linked coagulation disorder. The frequency of hemophilia A is 1-2 in 10,000 male births in all ethnic groups. About 50% of

**Nanjing BYabscience technology Co.,Ltd**



patients have severe hemophilia A with F8C activity less than 1% of normal; they have frequent spontaneous bleeding into joints, muscles and internal organs. Moderately severe hemophilia A occurs in about 10% of patients; F8C activity is 2-5% of normal, and there is bleeding after minor trauma. Mild hemophilia A, which occurs in 30-40% of patients, is associated with F8C activity of 5-30% and bleeding occurs only after significant trauma or surgery. Of particular interest for the understanding of the function of F8C is the category of CRM (cross-reacting material) positive patients (approximately 5%) that have considerable amount of F8C in their plasma (at least 30% of normal), but the protein is non-functional; i.e., the F8C activity is much less than the plasma protein level. CRM-reduced is another category of patients in which the F8C antigen and activity are reduced to approximately the same level. Most mutations are CRM negative, and probably affect the folding and stability of the protein.,domain:Domain F5/8 type C 2 is responsible for phospholipid-binding and essential for factor VIII activity.,function:Factor VIII, along with calcium and phospholipid, acts as a cofactor for factor IXa when it converts factor X to the activated form, factor Xa.,mass spectrometry:Disulfated PubMed:10368977,mass spectrometry:Monosulfated PubMed:10368977,mass spectrometry:Nonsulfated PubMed:10368977,mass spectrometry:Sulfated PubMed:10368977,mass spectrometry:Trisulfated PubMed:10368977,online information:Factor VIII entry,online information:Factor VIII mutation db,pharmaceutical:Available under the names Kogenate (Bayer) and Recombinate (Baxter and American Home Products). Used to treat hemophilia A.,PTM:Sulfation on Tyr-1699 is essential for binding vWF.,similarity:Belongs to the multicopper oxidase family.,similarity:Contains 1 F5/8 type C domain.,similarity:Contains 2 F5/8 type C domains.,similarity:Contains 3 F5/8 type A domains.,similarity:Contains 6 plastocyanin-like domains.,subunit:Interacts with vWF. vWF binding is essential for the stabilization of F8 in circulation.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658