



RGS16 (phospho Tyr168) Monoclonal Antibody

Catalog No BYmab-03535 Isotype IgG Reactivity Human;Mouse;Rat;Monkey Applications WB Gene Name RGS16 Protein Name Regulator of G-protein signaling 16 Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling and additional signaling and add		
Reactivity Human;Mouse;Rat;Monkey Applications WB Gene Name RGS16 Protein Name Regulator of G-protein signaling 16 Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; RRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by	Catalog No	BYmab-03535
Applications WB Gene Name RGS16 Protein Name Regulator of G-protein signaling 16 Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; RRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function: Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to Cys-2 and/or Cys-12. PTM-Phosphorylation on Tyn-168 upon EFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha., similarity. Contains 1 RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity lover levels of RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity contains 1 RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity contains 1 RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity contains 1 RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity contains 1 RGS domain. Issue specificity (AGP) activity on G(i)-alpha, similarity contains 1 RGS domain. Issue specificity Advancantly expressed in retina with lower levels of the results of the results are results and the results of the results and the results and the re	Isotype	IgG
Gene Name RGS16 Protein Name Regulator of G-protein signaling 16 Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade, PTM:Plaminotylated on Cys-2 and/of Cys-12, PTM:Phosphorylation on Tyr-168 uppersessed in retina with lower levels of RGS activity on G(i)-alpha, signaling in the phototransduction cascade, PTM:Plaminotylated on Cys-2 and/of Cys-12, PTM:Phosphorylation on Tyr-168 uppersessed in retina with lower levels of RGS domain, tissue specificity. Abundantly expressed in retina with lower levels of RGS domain, tissue specificity. Abundantly expressed in retina with lower levels of RGS domain, tissue specificity. Abundantly expressed in retina with lower levels of RGS domain.	Reactivity	Human;Mouse;Rat;Monkey
Protein Name Regulator of G-protein signaling 16 Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-audin form. Binds to G()-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12. PTM:Phosphorylation on Tyr-168 stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha, plane in retina with lower levels of ine tima with lower levels of ine ti	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G()-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12_PTM:Phosphorylation on Tyr-169 pone	Gene Name	RGS16
RGS16 around the phosphorylation site of Tyr168. AA range:141-190 Specificity Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function: Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulation the Cys-2 and/or Cys-12, PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha in retina with lower levels of RGS domain., tissue specificity: Abundantly expressed in retina with lower levels of in retina with lower levels of retina the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12, PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha in retina with lower levels of retina retina with lower levels of retina with low	Protein Name	Regulator of G-protein signaling 16
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade, PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha, similarity:Contains 1 RGS domain, itssue specificity:Abundantly expressed in retina with lower levels of very respected in retina with lower level	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction caded. PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha, similarity:Contains 1 RGS domain, tissue specificity:Abundantly expressed in retina with lower levels of in retina with lower levels of the phototransduction accelerating the kinetics of signaling in the phototransduction accelerating with lower levels of the phototransduction acceleration with lower levels of the phototransduction accelerating the kinetics of signal transduction acceleration acceleration with lower levels of the phototransduction acceleration w	Specificity	Phospho-RGS16 (Y168) Monoclonal Antibody detects endogenous levels of RGS16 protein only when phosphorylated at Y168.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha. But not to G(s)-alpha. May play a role in regulating the kinetics of signalling in the phototransduction on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP)-alpha. Simillarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of retina w	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12. PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha., similarity:Contains 1 RGS domain., tissue specificity:Abundantly expressed in retina with lower levels of	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signalling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12,.PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12., PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha., similarity:Contains 1 RGS domain., tissue specificity:Abundantly expressed in retina with lower levels of	Concentration	1 mg/ml
Synonyms RGS16; RGSR; Regulator of G-protein signaling 16; RGS16; A28-RGS14P; Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha, similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Purity	≥90%
Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein signaling Observed Band 30kD Cell Pathway Membrane; Lipid-anchor. Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade. PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Storage Stability	-20°C/1 year
Cell Pathway Membrane; Lipid-anchor. Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Synonyms	Retinal-specific RGS; RGS-r; hRGS-r; Retinally abundant regulator of G-protein
Tissue Specificity Abundantly expressed in retina with lower levels of expression in most other tissues. Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Observed Band	30kD
function: Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Cell Pathway	Membrane ; Lipid-anchor .
alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of	Tissue Specificity	
	Function	alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i)-alpha and G(o)-alpha, but not to G(s)-alpha. May play a role in regulating the kinetics of signaling in the phototransduction cascade.,PTM:Palmitoylated on Cys-2 and/or Cys-12.,PTM:Phosphorylation on Tyr-168 upon EGFR stimulation. Enhanced GTPase accelerating (GAP) activity on G(i)-alpha.,similarity:Contains 1 RGS domain.,tissue specificity:Abundantly expressed in retina with lower levels of

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	The protein encoded by this gene belongs to the 'regulator of G protein signaling' family. It inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits. It also may play a role in regulating the kinetics of signaling in the phototransduction cascade. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658