



MGST1 Monoclonal Antibody

Catalog No	BYmab-03955
Isotype	IgG
Reactivity	Human;Mouse;
Applications	WB
Gene Name	MGST1
Protein Name	Microsomal glutathione S-transferase 1
Immunogen	The antiserum was produced against synthesized peptide derived from human MGST1. AA range:42-91
Specificity	MGST1 Monoclonal Antibody detects endogenous levels of MGST1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MGST1; GST12; MGST; Microsomal glutathione S-transferase 1; Microsomal GST-1; Microsomal GST-I
Observed Band	18kD
Cell Pathway	Microsome . Mitochondrion outer membrane ; Peripheral membrane protein . Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	Highly expressed in liver.
Function	catalytic activity:RX + glutathione = HX + R-S-glutathione.,enzyme regulation:Activated by N-ethylmaleimide, except in the testis.,function:Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Has a wide substrate specificity.,similarity:Belongs to the MAPEG family.,subunit:Homotrimer.,tissue specificity:Highly expressed in liver.,
Background	The MAPEG (Membrane Associated Proteins in Eicosanoid and Glutathione metabolism) family consists of six human proteins, two of which are involved in the production of leukotrienes and prostaglandin E, important mediators of

Nanjing BYabscience technology Co.,Ltd



inflammation. Other family members, demonstrating glutathione S-transferase and peroxidase activities, are involved in cellular defense against toxic, carcinogenic, and pharmacologically active electrophilic compounds. This gene encodes a protein that catalyzes the conjugation of glutathione to electrophiles and the reduction of lipid hydroperoxides. This protein is localized to the endoplasmic reticulum and outer mitochondrial membrane where it is thought to protect these membranes from oxidative stress. Several transcript variants, some non-protein coding and some protein coding, have been found for this gene. [provided by RefSeq, May 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd