



# Tollip Monoclonal Antibody


<b>Catalog No</b>	BYmab-04248
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	TOLLIP
<b>Protein Name</b>	Toll-interacting protein
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human Tollip.
<b>Specificity</b>	Tollip Monoclonal Antibody detects endogenous levels of Tollip protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	TOLLIP; Toll-interacting protein
<b>Observed Band</b>	30kD
<b>Cell Pathway</b>	Cytoplasm .
<b>Tissue Specificity</b>	Brain,Eye,Placenta,Testis,
<b>Function</b>	function:Component of the signaling pathway of IL-1 and Toll-like receptors. Inhibits cell activation by microbial products. Recruits IRAK1 to the IL-1 receptor complex. Inhibits IRAK1 phosphorylation and kinase activity.,PTM:Phosphorylated by IRAK1 upon stimulation by IL-1 or microbial products.,similarity:Contains 1 CUE domain.,subunit:Oligomerizes. Binds to TLR2 and the TLR4-MD2 complex via its C-terminus. Exists as complex with IRAK1 in unstimulated cells. Upon IL-1 signaling, Tollip binds to the activated IL-1 receptor complex containing IL-1RI, IL-1RacP and the adapter protein MyD88, where it interacts with the TIR domain of IL-1RacP. MyD88 then triggers IRAK1 autophosphorylation, which in turn leads to the dissociation of IRAK1 from Tollip and IL-1RacP. Interacts with TOM1L2.,

**Nanjing BYabscience technology Co.,Ltd**



<b>Background</b>	This gene encodes a ubiquitin-binding protein that interacts with several Toll-like receptor (TLR) signaling cascade components. The encoded protein regulates inflammatory signaling and is involved in interleukin-1 receptor trafficking and in the turnover of IL1R-associated kinase. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2016],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using Tollip Monoclonal Antibody