



AXIN1 Monoclonal Antibody

Catalog No	BYmab-05368
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	AXIN1 AXIN
Protein Name	Axin-1 (Axis inhibition protein 1) (hAxin)
Immunogen	Synthesized peptide derived from human protein . at AA range: 190-270
Specificity	AXIN1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	94kD
Cell Pathway	Cytoplasm . Nucleus . Membrane . Cell membrane . MACF1 is required for its translocation to cell membrane (By similarity). On UV irradiation, translocates to the nucleus and colocalizes with DAAX (PubMed:17210684). .
Tissue Specificity	Ubiquitously expressed.
Function	disease:Defects in AXIN1 are involved in hepatocellular carcinoma (HCC) [MIM:114550].,disease:Hypermethylation of the AXIN1 promoter may be associated with caudal duplication anomaly [MIM:607864]. Caudal duplication anomaly is characterized by the occurrence of duplications of different organs in the caudal region.,function:Controls dorsoventral patterning via two opposing effects; down-regulates beta-catenin to inhibit the Wnt signaling pathway and ventralize embryos, but also dorsalizes embryos by activating a Wnt-independent JNK signaling pathway. In Wnt signaling, probably facilitates the phosphorylation of beta-catenin and APC by GSK3B. Likely to function as a tumor suppressor. Facilitates the phosphorylation of TP53 by HIPK2 upon ultraviolet irradiation. Wild-type axin 1 can induce apoptosis in hepatocellular and colorectal cancer

Nanjing BYabscience technology Co.,Ltd



cells. Enhances TGF-beta signaling by recruiting th

Background

This gene encodes a cytoplasmic protein which contains a regulation of G-protein signaling (RGS) domain and a dishevelled and axin (DIX) domain. The encoded protein interacts with adenomatosis polyposis coli, catenin beta-1, glycogen synthase kinase 3 beta, protein phosphate 2, and itself. This protein functions as a negative regulator of the wingless-type MMTV integration site family, member 1 (WNT) signaling pathway and can induce apoptosis. The crystal structure of a portion of this protein, alone and in a complex with other proteins, has been resolved. Mutations in this gene have been associated with hepatocellular carcinoma, hepatoblastomas, ovarian endometrioid adenocarcinomas, and medullablastomas. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd