



CBPA4 Monoclonal Antibody

| Catalog No | BYmab-05427 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | CPA4 CPA3 UNQ694/PRO1339 |
| Protein Name | Carboxypeptidase A4 (EC 3.4.17) (Carboxypeptidase A3) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | CBPA4 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 46kD |
| Cell Pathway | Secreted . |
| Tissue Specificity | Fetal expression in the adrenal gland, brain, heart, intestine, kidney, liver and lung. Except for fetal brain that shows no imprinting, expression was found preferentially from the maternal allele. |
| Function | cofactor:Binds 1 zinc ion per subunit.,function:Metalloprotease that could be involved in the histone hyperacetylation pathway.,induction:Up-regulated by inhibitors of histone dacetylation.,similarity:Belongs to the peptidase M14 family.,subunit:Interacts with LXN.,tissue specificity:Fetal expression in the adrenal gland, brain, heart, intestine, kidney, liver and lung. Except for fetal brain that shows no imprinting, expression was found preferentially from the maternal allele., |
| Background | This gene is a member of the carboxypeptidase A/B subfamily, and it is located in a cluster with three other family members on chromosome 7. Carboxypeptidases are zinc-containing exopeptidases that catalyze the release of |
| | |

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





| | carboxy-terminal amino acids, and are synthesized as zymogens that are activated by proteolytic cleavage. This gene could be involved in the histone hyperacetylation pathway. It is imprinted and may be a strong candidate gene for prostate cancer aggressiveness. [provided by RefSeq, Jul 2008], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Nanjing BYabscience technology Co.,Ltd