



# S6A19 Monoclonal Antibody

<b>Catalog No</b>	BYmab-06217
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	SLC6A19 B0AT1
<b>Protein Name</b>	Sodium-dependent neutral amino acid transporter B(0)AT1 (Solute carrier family 6 member 19) (System B(0) neutral amino acid transporter AT1)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	S6A19 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	69kD
<b>Cell Pathway</b>	Cell membrane ; Multi-pass membrane protein . Apical cell membrane ; Multi-pass membrane protein . Colocalizes with ACE2 on the apical membrane of cells lining villi of the jejunum, ileum and on kidney proximal tubules. .
<b>Tissue Specificity</b>	Robust expression in kidney and small intestine, with minimal expression in pancreas (PubMed:18424768, PubMed:15286787). Also expressed in stomach, liver, duodenum, ileocecum, colon and prostate. Not detected in testis, whole brain, cerebellum, fetal liver, spleen, skeletal muscle, uterus, heart or lung.
<b>Function</b>	disease:Defects in SLC6A19 are a cause of Hartnup disorder (HND) [MIM:234500]. HND is an autosomal recessive abnormality of renal and gastrointestinal neutral amino acid transport noted for its clinical variability. First described in 1956, HND is characterized by increases in the urinary and intestinal excretion of neutral amino acids. Individuals with typical Hartnup aminoaciduria may be asymptomatic, some develop a photosensitive pellagra-like rash, attacks of cerebellar ataxia and other neurological or psychiatric features. Although the definition of HND was originally based on clinical and biochemical abnormalities,

**Nanjing BYabscience technology Co.,Ltd**



its marked clinical heterogeneity has led to it being known as a disorder with a consistent pathognomonic neutral hyperaminoaciduria.,function:Transporter that mediates epithelial resorption of neutral amino acids across the apical membrane of epithelial cells in the kid

**Background**

This gene encodes a system B(0) transmembrane protein that actively transports most neutral amino acids across the apical membrane of epithelial cells. Mutations in this gene result in Hartnup disorder. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658