



## TBG2 mouse mAb

Catalog No	BYmab-08498
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	TUBG2
Protein Name	TBG2
Immunogen	Synthesized peptide derived from human TBG2 AA range: 113-163
Specificity	This antibody detects endogenous levels of TBG2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.
Tissue Specificity	
Function	function: Tubulin is the major constituent of microtubules. Gamma tubulin is found at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of microtubule assembly., similarity: Belongs to the tubulin family.,
Function  Background	at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of
	at microtubule organizing centers (MTOC) such as the spindle poles or the centrosome, suggesting that it is involved in the minus-end nucleation of

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using TBG2 mouse mAb