



## SLAP2 mouse mAb

Catalog No	BYmab-10081
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SLA2 C20orf156 SLAP2
Protein Name	Src-like-adapter 2 (Modulator of antigen receptor signaling) (MARS) (Src-like adapter protein 2) (SLAP-2)
Immunogen	Synthesized peptide derived from human SLAP2 AA range: 159-209
Specificity	This antibody detects endogenous levels of human SLAP2
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.;
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd