



NO66 mouse mAb

Catalog No	BYmab-11227
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	NO66 C14orf169 MAPJD
Protein Name	NO66
Immunogen	Synthesized peptide derived from human NO66 AA range: 533-583
Specificity	This antibody detects endogenous levels of NO66 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Nucleus, nucleolus. Nucleus, nucleoplasm. Granular part of nucleoli.
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Nucleus, nucleolus. Nucleus, nucleoplasm. Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them associated with nucleoli. Widely expressed. Overexpressed in lung carcinomas. function:May play a role in ribosome biogenesis and in the replication or remodeling of certain heterochromatic region.,similarity:Belongs to the MINA53/NO66 family.,similarity:Contains 1 JmjC domain.,subcellular
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Nucleus, nucleolus. Nucleus, nucleoplasm. Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them associated with nucleoli. Widely expressed. Overexpressed in lung carcinomas. function:May play a role in ribosome biogenesis and in the replication or remodeling of certain heterochromatic region.,similarity:Belongs to the MINA53/NO66 family.,similarity:Contains 1 JmjC domain.,subcellular location:Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Nucleus, nucleolus. Nucleus, nucleoplasm. Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them associated with nucleoli. Widely expressed. Overexpressed in lung carcinomas. function:May play a role in ribosome biogenesis and in the replication or remodeling of certain heterochromatic region.,similarity:Belongs to the MINA53/NO66 family.,similarity:Contains 1 JmjC domain.,subcellular location:Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NO66 mouse mAb