



PATZ1 mouse mAb

Catalog No	BYmab-11236
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	PATZ1 PATZ RIAZ ZBTB19 ZNF278 ZSG
Protein Name	PATZ1
Immunogen	Synthesized peptide derived from human PATZ1 AA range: 535-585
Specificity	This antibody detects endogenous levels of PATZ1 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitous.
Function	disease:A chromosomal aberration involving PATZ1 is associated with small round cell sarcoma. Translocation t(1;22)(p36.1;q12) with EWSR1.,function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 A.T hook DNA-binding domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 7 C2H2-type zinc fingers.,tissue specificity:Ubiquitous.,
Background	The protein encoded by this gene contains an A-T hook DNA binding motif which usually binds to other DNA binding structures to play an important role in chromatin modeling and transcription regulation. Its Poz domain is thought to function as a site for protein-protein interaction and is required for transcriptional repression, and the zinc-fingers comprise the DNA binding domain. Since the

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



encoded protein has typical features of a transcription factor, it is postulated to be
a repressor of gene expression. In small round cell sarcoma, this gene is fused to
EWS by a small inversion of 22q, then the hybrid is thought to be translocated
(t(1;22)(p36.1;q12). The rearrangement of chromosome 22 involves intron 8 of
ÈWS and exon 1 of this gene creating a chimeric sequence containing the
transactivation domain of EWS fused to zinc finger domain of this protein. This is a
distinct example of an intra-chromosomal rearrangement of chromosome 22. Four
alternatively spliced transcript variants are described for this gene. [provided by
RefSeg, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

