



CAR Monoclonal Antibody


Catalog No	BYmab-13161
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CXADR
Protein Name	Coxsackievirus and adenovirus receptor
Immunogen	The antiserum was produced against synthesized peptide derived from human CXADR. AA range:1-50
Specificity	CAR Monoclonal Antibody detects endogenous levels of CAR protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CXADR; CAR; Coxsackievirus and adenovirus receptor; CAR; hCAR; CVB3-binding protein; Coxsackievirus B-adenovirus receptor; HCVADR
Observed Band	40kD
Cell Pathway	[Isoform 1]: Cell membrane ; Single-pass type I membrane protein . Basolateral cell membrane ; Single-pass type I membrane protein . Cell junction, tight junction . Cell junction, adherens junction . In epithelial cells localizes to the apical junction complex composed of tight and adherens junctions (PubMed:12297051). In airway epithelial cells localized to basolateral membrane but not to apical surface (PubMed:11316797). .; [Isoform 3]: Secreted .; [Isoform 4]: Secreted .; [Isoform 5]: Secreted .
Tissue Specificity	Expressed in pancreas, brain, heart, small intestine, testis, prostate and at a lower level in liver and lung. Isoform 5 is ubiquitously expressed. Isoform 3 is expressed in heart, lung and pancreas. In skeletal muscle, isoform 1 is found at the neuromuscular junction and isoform 2 is found in blood vessels. In cardiac muscle, isoform 1 and isoform 2 are found at intercalated disks. In heart expressed in subendothelial layers of the vessel wall but not in the luminal endothelial surface. Expression is elevated in hearts with dilated cardiomyopathy.

Nanjing BYabscience technology Co.,Ltd



Function	domain:The Ig-like C2-type 1 domain probably mediates homodimerization and interaction with JAML.,domain:The PDZ-binding motif mediates interaction with MPDZ and BAIAP1.,function:Component of the epithelial apical junction complex that is essential for the tight junction integrity. Proposed to function as a homophilic cell adhesion molecule. Recruits MPDZ to intercellular contact sites. Probably involved in transepithelial migration of polymorphonuclear leukocytes (PMN) through adhesive interactions with AMICA1/JAML located in the plasma membrane of PMN.,PTM:N-glycosylated.,PTM:Palmitoylated on Cys-259 and/or Cys-260; required for proper localization to the plasma membrane.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subcellular location:In epithelial cells localizes to the apical junction complex composed of tight and adherens junctions. In airway epithelial ce
Background	The protein encoded by this gene is a type I membrane receptor for group B coxsackieviruses and subgroup C adenoviruses. Several transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene are found on chromosomes 15, 18, and 21. [provided by RefSeq, May 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CAR Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658