



Olfactory receptor 4F6 Monoclonal Antibody

Catalog No BYmab-13534 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name OR4F6 Protein Name Olifactory receptor 4F6 Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor 0R15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Image: Specificity Function function: Odorant receptor family 4 subfamily F member 6(OR4F6) Homo sapiens olfactory re		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR4F6 Protein Name Olfactory receptor 4F6 Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors of a large family of G-protein-coupled receptors (GPCR) arising from single coding-evon queen. Olfactory receptors (GPCR) arising from single coding-evon queen.	Catalog No	BYmab-13534
Applications WB Gene Name OR4F6 Protein Name Olfactory receptor 4F6 Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a	Isotype	IgG
Gene Name OR4F6 Protein Name Olfactory receptor 4F6 Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 4F6 Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors share a series of a large family of G-protein-coupled receptors share a colfactory receptor share a colfactory receptor share a colfactory receptor share a colfacto	Gene Name	OR4F6
OR4F6. AA range:55-104 Specificity Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptor (GPCR) arising from single coding-exon genes. Olfactor-ceceptors (GPCR) arising from single coding-exon genes. Olfactor-ceceptor sceptors (GPCR)	Protein Name	Olfactory receptor 4F6
Olfactory receptor 4F6 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor ,,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Specificity	Olfactory receptor 4F6 Monoclonal Antibody detects endogenous levels of Olfactory receptor 4F6 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors grising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor 0R15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Concentration	1 mg/ml
Synonyms OR4F6; OR4F12; Olfactory receptor 4F6; Olfactory receptor 4F12; Olfactory receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Purity	≥90%
receptor OR15-15 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Synonyms	
Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Observed Band	35kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a		
Background Olfactory receptor family 4 subfamily F member 6(OR4F6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a	Cell Pathway	
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a		
	Tissue Specificity	Cell membrane; Multi-pass membrane protein. function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Olfactory receptor 4F6 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658