



p47-phox Polyclonal Antibody

Catalog No	BYab-19137
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	NCF1
Protein Name	Neutrophil cytosol factor 1
Immunogen	The antiserum was produced against synthesized peptide derived from human p47 phox. AA range:341-390
Specificity	p47-phox Polyclonal Antibody detects endogenous levels of p47-phox protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NCF1; NOXO2; SH3PXD1A; Neutrophil cytosol factor 1; NCF-1; 47 kDa autosomal chronic granulomatous disease protein; 47 kDa neutrophil oxidase factor; NCF-47K; Neutrophil NADPH oxidase factor 1; Nox organizer 2; Nox-organizing protein 2; SH3
Observed Band	44kD
Cell Pathway	Cytoplasm, cytosol . Membrane ; Peripheral membrane protein ; Cytoplasmic side
Tissue Specificity	Detected in peripheral blood monocytes and neutrophils (at protein level).
Function	disease:Defects in NCF1 are the cause of chronic granulomatous disease autosomal recessive cytochrome-b-positive type 1 (CGD1) [MIM:233700]. Chronic granulomatous disease is a genetically heterogeneous disorder characterized by the inability of neutrophils and phagocytes to kill microbes that they have ingested. Patients suffer from life-threatening bacterial/fungal infections.,function:NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide

Nanjing BYabscience technology Co.,Ltd



production).,online information:NCF1 deficiency database,similarity:Contains 1 PX (phox homology) domain.,similarity:Contains 2 SH3 domains.,subunit:Interacts with NOXA1.,

Background

The protein encoded by this gene is a 47 kDa cytosolic subunit of neutrophil NADPH oxidase. This oxidase is a multicomponent enzyme that is activated to produce superoxide anion. Mutations in this gene have been associated with chronic granulomatous disease. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images




Western Blot analysis of various cells using p47-phox Polyclonal Antibody




Western Blot analysis of COLO205 cells using p47-phox Polyclonal Antibody


Nanjing BYabs science technology Co.,Ltd



Immunohistochemistry analysis of paraffin-embedded human tonsil tissue, using p47 phox Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HT-29, COLO205, and HepG2 cells, using p47 phox Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using p47 phox antibody.