



PFAAP5 Polyclonal Antibody

Catalog No	BYab-19152
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA;IHC
Gene Name	N4BP2L2
Protein Name	NEDD4-binding protein 2-like 2
Immunogen	The antiserum was produced against synthesized peptide derived from human N4BP2L2. AA range:358-407
Specificity	PFAAP5 Polyclonal Antibody detects endogenous levels of PFAAP5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	N4BP2L2; CG005; PFAAP5; NEDD4-binding protein 2-like 2; Phosphonoformate immuno-associated protein 5
Observed Band	67kD
Cell Pathway	nucleus,transcriptional repressor complex,extracellular exosome,
Tissue Specificity	Colon,Thalamus,
Function	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,
Background	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,
matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998


监督电话: 15950492658




Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of LOVO cells using PFAAP5 Polyclonal Antibody




Western blot analysis of lysates from LOVO cells, using N4BP2L2 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabs science technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Western blot analysis of the lysates from K562 cells using N4BP2L2 antibody.



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).