



17 β -HSD4 Polyclonal Antibody

Catalog No	BYab-19269
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	HSD17B4
Protein Name	Peroxisomal multifunctional enzyme type 2
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human HSD17B4. AA range:41-90
Specificity	17 β -HSD4 Polyclonal Antibody detects endogenous levels of 17 β -HSD4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HSD17B4; EDH17B4; Peroxisomal multifunctional enzyme type 2; MFE-2; 17-beta-hydroxysteroid dehydrogenase 4; 17-beta-HSD 4; D-bifunctional protein; DBP; Multifunctional protein 2; MPF-2
Observed Band	80kD
Cell Pathway	Peroxisome .
Tissue Specificity	Present in many tissues with highest concentrations in liver, heart, prostate and testis.
Function	catalytic activity:(24R,25R)-3-alpha,7-alpha,12-alpha,24-tetrahydroxy-5-beta-cholestanoyl-CoA = (24E)-3-alpha,7-alpha,12-alpha-trihydroxy-5-beta-cholest-24-enoyl-CoA + H(2)O.,catalytic activity:(S)-3-hydroxyacyl-CoA + NAD(+) = 3-oxoacyl-CoA + NADH.,disease:Defects in HSD17B4 are a cause of D-bifunctional protein deficiency (DBPD) [MIM:261515]. DBPD is a disorder of peroxisomal fatty acid beta-oxidation.,function:Bifunctional enzyme acting on the peroxisomal beta-oxidation pathway for fatty acids. Catalyzes the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty


Nanjing BYabscience technology Co.,Ltd




	acids.,pathway:Lipid metabolism; fatty acid beta-oxidation.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Contains 1 SCP2 domain.,tissue specificity:Present in many tissues with highest concentrations in liver, heart, prostate and testis.,
Background	hydroxysteroid 17-beta dehydrogenase 4(HSD17B4) Homo sapiens The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of HuvEc, K562 cells using 17 β -HSD4 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

Nanjing BYabs science technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



Western blot analysis of lysate from K562, HUVEC cells, using HSD17B4 Antibody.