



PXR Polyclonal Antibody

Catalog No	BYab-19682
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	NR1I2
Protein Name	Nuclear receptor subfamily 1 group I member 2
Immunogen	The antiserum was produced against synthesized peptide derived from human NR1I2. AA range:91-140
Specificity	PXR Polyclonal Antibody detects endogenous levels of PXR protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NR1I2; PXR; Nuclear receptor subfamily 1 group I member 2; Orphan nuclear receptor PAR1; Orphan nuclear receptor PXR; Pregnane X receptor; Steroid and xenobiotic receptor; SXR
Observed Band	50kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in liver, colon and small intestine.
Function	function:Orphan receptor; its natural ligand is probably pregnane. Binds to a response element in the CYP3A4 and ABCB1/MDR1 genes promoter. Activates its expression in response to a wide variety of endobiotics and xenobiotics.,induction:Activated by naturally occurring steroids such as pregnenolone and progesterone.,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Forms a heterodimer with RXR.,tissue specificity:Expressed in liver, colon and small intestine.,

Nanjing BYabscience technology Co.,Ltd



Background

This gene product belongs to the nuclear receptor superfamily, members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Zhao, Guo-Ping, et al. "Imidacloprid increases intestinal permeability by disrupting tight junctions." *Ecotoxicology and Environmental Safety* 222 (2021): 112476.