



tPA Polyclonal Antibody

Catalog No	BYab-20571
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	PLAT
Protein Name	Tissue-type plasminogen activator
Immunogen	The antiserum was produced against synthesized peptide derived from human tPA. AA range:38-87
Specificity	tPA Polyclonal Antibody detects endogenous levels of tPA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PLAT; Tissue-type plasminogen activator; t-PA; t-plasminogen activator; tPA; Alteplase; Reteplase
Observed Band	63kD
Cell Pathway	Secreted, extracellular space.
Tissue Specificity	Synthesized in numerous tissues (including tumors) and secreted into most extracellular body fluids, such as plasma, uterine fluid, saliva, gingival crevicular fluid, tears, seminal fluid, and milk.
Function	catalytic activity:Specific cleavage of Arg- -Val bond in plasminogen to form plasmin.,disease:Increased activity of TPA is the cause of hyperfibrinolysis [MIM:173370]. Hyperfibrinolysis leads to excessive bleeding. Defective release of TPA causes hypofibrinolysis, leading to thrombosis or embolism.,domain:Both FN1 and EGF-like domains are important for binding to LRP1.,domain:Both FN1 and one of the kringle domains are required for binding to fibrin.,domain:The FN1 domain mediates binding to annexin A2.,domain:The second kringle domain is implicated in binding to cytokeratin-8 and to the endothelial cell surface binding site.,function:Converts the abundant, but inactive, zymogen plasminogen to

Nanjing BYabscience technology Co.,Ltd



plasmin by hydrolyzing a single Arg-Val bond in plasminogen. By controlling plasmin-mediated proteolysis, it plays an important role in tissue remodeling and degradation, in cell migration and man

Background

This gene encodes tissue-type plasminogen activator, a secreted serine protease that converts the proenzyme plasminogen to plasmin, a fibrinolytic enzyme. The encoded preproprotein is proteolytically processed by plasmin or trypsin to generate heavy and light chains. These chains associate via disulfide linkages to form the heterodimeric enzyme. This enzyme plays a role in cell migration and tissue remodeling. Increased enzymatic activity causes hyperfibrinolysis, which manifests as excessive bleeding, while decreased activity leads to hypofibrinolysis, which can result in thrombosis or embolism. Alternative splicing of this gene results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998


监督电话: 15950492658



Products Images



Western Blot analysis of A549 cells using tPA Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from A549 cells, using tPA antibody.

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658