



PRX I Polyclonal Antibody

Catalog No	BYab-20662
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	PRDX1
Protein Name	Peroxiredoxin-1
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human PRDX1. AA range:31-80
Specificity	PRX I Polyclonal Antibody detects endogenous levels of PRX I protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PRDX1; PAGA; PAGB; TDPX2; Peroxiredoxin-1; Natural killer cell-enhancing factor A; NKEF-A; Proliferation-associated gene protein; PAG; Thioredoxin peroxidase 2; Thioredoxin-dependent peroxide reductase 2
Observed Band	21kD
Cell Pathway	Cytoplasm . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Tissue Specificity	Brain,Cajal-Retzius cell,Fetal brain cortex,Urinary bladder,
Function	catalytic activity:2 R'-SH + ROOH = R'-S-S-R' + H(2)O + ROH.,function:Involved in redox regulation of the cell. Reduces peroxides with reducing equivalents provided through the thioredoxin system but not from glutaredoxin. May play an important role in eliminating peroxides generated during metabolism. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).,induction:Constitutively expressed in most human cells; is induced to higher levels upon serum stimulation in untransformed and transformed cells.,miscellaneous:Inactivated upon oxidative stress by overoxidation of Cys-52

Nanjing BYabs science technology Co.,Ltd



to Cys-SO(2)H and Cys-SO(3)H. Cys-SO(2)H is retroreduced to Cys-SOH after removal of H₂O₂, while Cys-SO(3)H may be irreversibly oxidized. miscellaneous: The active site is the redox-active Cys-52 oxidized to Cys-SOH.

Background

This gene encodes a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen peroxide and alkyl hydroperoxides. The encoded protein may play an antioxidant protective role in cells, and may contribute to the antiviral activity of CD8(+) T-cells. This protein may have a proliferative effect and play a role in cancer development or progression. Four transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq, Jan 2011],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of HeLa cells using PRX I Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Western blot analysis of lysate from HeLa cells, using PRDX1 Antibody.