



RO52 Polyclonal Antibody

Catalog No	BYab-21463
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	TRIM21 RNF81 RO52 SSA1
Protein Name	E3 ubiquitin-protein ligase TRIM21 (EC 6.3.2.-) (52 kDa Ro protein) (52 kDa ribonucleoprotein autoantigen Ro/SS-A) (RING finger protein 81) (Ro(SS-A)) (Sjogren syndrome type A antigen) (SS-A) (Tripar
Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Specificity	RO52 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	52kD
Cell Pathway	Cytoplasm . Cytoplasmic vesicle, autophagosome . Nucleus . Cytoplasm, P-body . Enters the nucleus upon exposure to nitric oxide. Localizes to small dot- or rod-like structures in the cytoplasm, called processing bodies (P-bodies) that are located underneath the plasma membrane and also diffusely in the cytoplasm. They are located along the microtubules and are highly motile in cells. Colocalizes with DCP2 in P-bodies.
Tissue Specificity	Isoform 1 and isoform 2 are expressed in fetal and adult heart and fetal lung.
Function	disease:Sera from patients with systemic lupus erythematosus (SLE) or primary Sjogren syndrome often contain antibodies that react with normal cellular RO protein as if this antigen was foreign..function:Ribonucleoprotein particle composed of a single polypeptide and one of four small RNA molecules. It is present in all mammalian cells studied but has no known function. At least two isoforms are present in nucleated and red blood cells, and tissue specific

Nanjing BYabs science technology Co.,Ltd



differences in RO/SSA proteins have been identified. The common feature of these proteins is their ability to bind HY RNAs.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,

Background

This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The encoded protein is part of the RoSSA ribonucleoprotein, which includes a single polypeptide and one of four small RNA molecules. The RoSSA particle localizes to both the cytoplasm and the nucleus. RoSSA interacts with autoantigens in patients with Sjogren syndrome and systemic lupus erythematosus. Alternatively spliced transcript variants for this gene have been described but the full-length nature of only one has been determined. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658