



ATS14 Polyclonal Antibody

Catalog No	BYab-21506
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	ADAMTS14
Protein Name	A disintegrin and metalloproteinase with thrombospondin motifs 14 (ADAM-TS 14) (ADAM-TS14) (ADAMTS-14) (EC 3.4.24.-)
Immunogen	Synthesized peptide derived from human protein . at AA range: 190-270
Specificity	ATS14 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	134kD
Cell Pathway	Secreted, extracellular space, extracellular matrix .
Tissue Specificity	Expressed in retina and at low levels in brain, lung and placenta (PubMed:11779638). High expression in fetal tissues (PubMed:11867212).
Function	domain:The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix.,function:Has a aminoprocollagen type I activity processing activity in the absence of ADAMTS2. Seems to be synthesized as a latent enzyme that requires activation to display aminoprocollagen peptidase activity.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Contains 1 disintegrin domain.,similarity:Contains 1 peptidase M12B domain.,similarity:Contains 1 PLAC domain.,similarity:Contains 4 TSP type-1 domains.,tissue specificity:Expressed in retina and at low levels in brain, lung and placenta. High expression in fetal tissues.,
Background	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the

Nanjing BYabscience technology Co.,Ltd



family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature enzyme. This enzyme cleaves amino-terminal propeptides from type I procollagen, a necessary step in the formation of collagen fibers. Mutations in this gene may be associated with osteoarthritis in human patients. [provided by RefSeq, May 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658