



PAIP1 Polyclonal Antibody

Catalog No	BYab-22162
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PAIP1
Protein Name	Polyadenylate-binding protein-interacting protein 1 (PABP-interacting protein 1) (PAIP-1) (Poly(A)-binding protein-interacting protein 1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 100-180
Specificity	PAIP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	52kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Kidney,Placenta,Skin,Uterus,
Function	caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,domain:Only the PABPC1-interacting motif-1 (PAM1) stimulates translation initiation.,function:Acts as a coactivator in the regulation of translation initiation of poly(A)-containing mRNAs. Its stimulatory activity on translation is mediated via its action on PABPC1. Competes with PAIP2 for binding to PABPC1. Its association with EIF4A and PABPC1 may potentiate contacts between mRNA termini. May also be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain.,similarity:Contains 1 MIF4G domain.,subunit:Interacts with the RRM1-RRM2 and C-terminus regions of PABPC1

Nanjing BYabscience technology Co.,Ltd



Background	The protein encoded by this gene interacts with poly(A)-binding protein and with the cap-binding complex eIF4A. It is involved in translational initiation and protein biosynthesis. Overexpression of this gene in COS7 cells stimulates translation. Alternative splicing occurs at this locus and three transcript variants encoding three distinct isoforms have been identified. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images