



RPGF6 Polyclonal Antibody

Catalog No	BYab-22235
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	RAPGEF6 PDZGEF2
Protein Name	Rap guanine nucleotide exchange factor 6 (PDZ domain-containing guanine nucleotide exchange factor 2) (PDZ-GEF2) (RA-GEF-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 460-540
Specificity	RPGF6 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	176kD
Cell Pathway	Cytoplasm . Cell membrane . Upon binding to M-Ras, it translocates to the plasma membrane.
Tissue Specificity	Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.
Function	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,

Nanjing BYabscience technology Co.,Ltd



Background	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images