



TRPV6 Polyclonal Antibody

Catalog No	BYab-22528
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	TRPV6 ECAC2
Protein Name	Transient receptor potential cation channel subfamily V member 6 (TrpV6) (CaT-like) (CaT-L) (Calcium transport protein 1) (CaT1) (Epithelial calcium channel 2) (ECaC2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TRPV6 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	79kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Expressed at high levels in the gastrointestinal tract, including esophagus, stomach, duodenum, jejunum, ileum and colon, and in pancreas, placenta, prostate and salivary gland. Expressed at moderate levels in liver, kidney and testis. Expressed in trophoblasts of placenta villus trees (at protein level)(PubMed:23612980). Expressed in locally advanced prostate cancer, metastatic and androgen-insensitive prostatic lesions but not detected in healthy prostate tissue and benign prostatic hyperplasia.
Function	function:Calcium selective cation channel probably involved in Ca(2+) uptake in various tissues, including Ca(2+) reabsorption in intestine. The channel is activated by low internal calcium level, probably including intracellular calcium store depletion, and the current exhibits an inward rectification. Inactivation includes both, a rapid Ca(2+)-dependent and a slower

Nanjing BYabscience technology Co.,Ltd



Ca(2+)-calmodulin-dependent mechanism, the latter may be regulated by phosphorylation. In vitro, is slowly inhibited by Mg(2+) in a voltage-independent manner. Heteromeric assembly with TRPV5 seems to modify channel properties. TRPV5-TRPV6 heteromultimeric concatemers exhibit voltage-dependent gating.,online information:TRPV6 entry,PTM:Glycosylated.,PTM:Phosphorylation by PRKCA at the calmodulin binding site delays channel inactivation.,similarity:Belongs to the transient receptor family. TrpV subfamily.,similarity:Contains

Background

transient receptor potential cation channel subfamily V member 6 (TRPV6) Homo sapiens This gene encodes a member of a family of multipass membrane proteins that functions as calcium channels. The encoded protein contains N-terminal ankyrin repeats, which are required for channel assembly and regulation. Translation initiation for this protein occurs at a non-AUG start codon that is decoded as methionine. This gene is situated next to a closely related gene for transient receptor potential cation channel subfamily V member 5 (TRPV5). This locus has experienced positive selection in non-African populations, resulting in several non-synonymous codon differences among individuals of different genetic backgrounds. [provided by RefSeq, Feb 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658