



INT6 Polyclonal Antibody

Catalog No	BYab-22800
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	INTS6 DBI1 DDX26 DDX26A
Protein Name	Integrator complex subunit 6 (Int6) (DBI-1) (Protein DDX26) (Protein deleted in cancer 1) (DICE1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 550-630
Specificity	INT6 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	97kD
Cell Pathway	Nucleus .
Tissue Specificity	Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.
Function	function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes. May have a tumor suppressor role; an ectopic expression suppressing tumor cell growth.,induction:Frequently down-regulated in nonsmall cell lung carcinomas and prostate cancers. Down-regulation in prostate cancer is due to CpG hypermethylation of its promoter. However, some involvement in cancer is unclear.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the Integrator subunit 6 family.,similarity:Contains 1 VWFA domain.,subunit:Belongs to the multiprotein complex Integrator, at least

Nanjing BYabscience technology Co.,Ltd



composed of INTS1, INTS2, INTS3, INTS4,

Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. The protein encoded by this gene is a DEAD box protein that is part of a complex that interacts with the C-terminus of RNA polymerase II and is involved in 3' end processing of snRNAs. In addition, this gene is a candidate tumor suppressor and is located in the critical region of loss of heterozygosity (LOH). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658