



SYCP1 Polyclonal Antibody

Catalog No	BYab-22883
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	SYCP1 SCP1
Protein Name	Synaptonemal complex protein 1 (SCP-1) (Cancer/testis antigen 8) (CT8)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SYCP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	107kD
Cell Pathway	Nucleus . Chromosome . Chromosome, centromere . In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex (By similarity). Only rarely detected at centromeres during leptotene and zygotene. Detected at centromeres during mid-diplotene, when it is no longer present along chromosome arms. No longer detected at centromeres at later stages of meiosis (By similarity). .
Tissue Specificity	Testis.
Function	domain:Consists of an alpha-helical stretch of 700 AA residues, flanked by N- and C-terminal globular domains. The C-terminal domain has DNA-binding capacity.,function:Major component of the transverse filaments of synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase.,subcellular location:In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Found only where the chromosome

Nanjing BYabs science technology Co.,Ltd



	cores are synapsed. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex.,subunit:Found in a complex with SYCE1 and SYCE2. Interacts with SYCE1 and SYCE2.,tissue specificity:Testis.,
Background	domain:Consists of an alpha-helical stretch of 700 AA residues, flanked by N- and C-terminal globular domains. The C-terminal domain has DNA-binding capacity.,function:Major component of the transverse filaments of synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase.,subcellular location:In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Found only where the chromosome cores are synapsed. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex.,subunit:Found in a complex with SYCE1 and SYCE2. Interacts with SYCE1 and SYCE2.,tissue specificity:Testis.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658