



NCPR Polyclonal Antibody

Catalog No	BYab-22911
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	POR CYPOR
Protein Name	NADPH--cytochrome P450 reductase (CPR) (P450R) (EC 1.6.2.4)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NCPR Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	74kD
Cell Pathway	Endoplasmic reticulum membrane ; Single-pass membrane protein ; Cytoplasmic side .
Tissue Specificity	Aorta endothelial cell,Liver,Lung,Small intestine,
Function	catalytic activity:NADPH + n oxidized hemoprotein = NADP(+) + n reduced hemoprotein.,cofactor:FAD.,cofactor:FMN.,disease:Defects in POR are a cause of isolated disordered steroidogenesis (IDS) [MIM:201750].,disease:Defects in POR are the cause of adrenal hyperplasia variant type (AHV) [MIM:201750]; also known as Antley-Bixler syndrome-like phenotype with disordered steroidogenesis. AHV is a rare variant of congenital adrenal hyperplasia. It is an autosomal recessive disorder with apparent combined P450C17 and P450C21 deficiency. Affected girls are born with ambiguous genitalia, but their circulating androgens are low and virilization does not progress. Conversely, affected boys are sometimes born undermasculinized. Boys and girls can also present with bone

Nanjing BYabscience technology Co.,Ltd



malformations, in some cases resembling the pattern seen in patients with Antley-Bixler syndrome.,function:This enzyme is required fo

Background

This gene encodes an endoplasmic reticulum membrane oxidoreductase with an FAD-binding domain and a flavodoxin-like domain. The protein binds two cofactors, FAD and FMN, which allow it to donate electrons directly from NADPH to all microsomal P450 enzymes. Mutations in this gene have been associated with various diseases, including apparent combined P450C17 and P450C21 deficiency, amenorrhea and disordered steroidogenesis, congenital adrenal hyperplasia and Antley-Bixler syndrome. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cell lysis. Primary Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS23920 was diluted at 1:10000



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).