



HUWE1 Polyclonal Antibody

Catalog No	BYab-22917
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF
Gene Name	HUWE1 KIAA0312 KIAA1578 UREB1 HSPC272
Protein Name	E3 ubiquitin-protein ligase HUWE1 (EC 6.3.2.-) (ARF-binding protein 1) (ARF-BP1) (HECT, UBA and WWE domain-containing protein 1) (Homologous to E6AP carboxyl terminus homologous protein 9) (HectH9) (L
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HUWE1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-300. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	481kD
Cell Pathway	Cytoplasm . Nucleus . Mitochondrion . Mainly expressed in the cytoplasm of most tissues, except in the nucleus of spermatogonia, primary spermatocytes and neuronal cells (By similarity). Recruited to mitochondria following interaction with AMBRA1 (PubMed:30217973). .
Tissue Specificity	Weakly expressed in heart, brain and placenta but not in other tissues. Expressed in a number of cell lines, predominantly in those from colorectal carcinomas.
Function	disease:A chromosomal microduplication involving HUWE1 and HSD17B10 is the cause of mental retardation X-linked type 17 (MRX17) [MIM:300705]; also known as mental retardation X-linked type 31 (MRX31). Mental retardation is characterized by significantly sub-average general intellectual functioning associated with impairments in adaptative behavior and manifested during the developmental period. In contrast to syndromic or specific X-linked mental retardation which also present with associated physical, neurological and/or psychiatric manifestations, intellectual deficiency is the only primary symptom of

Nanjing BYabscience technology Co.,Ltd



non-syndromic X-linked mental retardation.,disease:Defects in HUWE1 are the cause of mental retardation syndromic X-linked Turner type (MRXST) [MIM:300706]; also known as mental retardation and macrocephaly syndrome. MRXST shows clinical variability. Associated phenotypes include macrocep

Background

This gene encodes a protein containing a C-terminal HECT (E6AP type E3 ubiquitin protein ligase) domain that functions as an E3 ubiquitin ligase. The encoded protein is required for the ubiquitination and subsequent degradation of the anti-apoptotic protein Mcl1 (myeloid cell leukemia sequence 1 (BCL2-related)). This protein also ubiquitinates the p53 tumor suppressor, core histones, and DNA polymerase beta. Mutations in this gene are associated with Turner type X-linked syndromic mental retardation. [provided by RefSeq, Aug 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human brain tumor. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).