



PKN3 Polyclonal Antibody

Catalog No	BYab-22952
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PKN3 PKNBETA
Protein Name	Serine/threonine-protein kinase N3 (EC 2.7.11.13) (Protein kinase PKN-beta) (Protein-kinase C-related kinase 3)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PKN3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	97kD
Cell Pathway	Nucleus . Cytoplasm, perinuclear region . Nuclear and perinuclear Golgi region.
Tissue Specificity	Expressed in prostate tumors and various cancer cell lines. Not expressed in adult tissues.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,domain:The C1 domain does not bind the diacylglycerol (DAG).,enzyme regulation:Two specific sites, Thr-718 (activation loop of the kinase domain) and Thr-860 (turn motif), need to be phosphorylated for its full activation.,function:Contributes to invasiveness in malignant prostate cancer.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 3 REM (Hr1) repeats.,subcellular location:Nuclear and perinuclear Golgi region.,tissue specificity:Expressed in prostate tumors and various cancer cell lines. Not

Nanjing BYabscience technology Co.,Ltd



expressed in adult tissues.,

Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,domain:The C1 domain does not bind the diacylglycerol (DAG).,enzyme regulation:Two specific sites, Thr-718 (activation loop of the kinase domain) and Thr-860 (turn motif), need to be phosphorylated for its full activation.,function:Contributes to invasiveness in malignant prostate cancer.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 3 REM (Hr1) repeats.,subcellular location:Nuclear and perinuclear Golgi region.,tissue specificity:Expressed in prostate tumors and various cancer cell lines. Not expressed in adult tissues.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658