



SLPI Polyclonal Antibody

Catalog No	BYab-23192
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	SLPI WAP4 WFDC4
Protein Name	Antileukoproteinase (ALP) (BLPI) (HUSI-1) (Mucus proteinase inhibitor) (MPI) (Protease inhibitor WAP4) (Secretory leukocyte protease inhibitor) (Seminal proteinase inhibitor) (WAP four-disulfide core)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SLPI Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	14kD
Cell Pathway	Secreted .
Tissue Specificity	Detected in blood plasma (PubMed:24352879). Detected in bone marrow myeloid cells (PubMed:24352879). Detected in airway sputum (PubMed:2039600). Detected in parotid gland secretions (PubMed:3462719). Detected in seminal plasma (at protein level) (PubMed:3485543). Detected in uterus cervix (PubMed:3533531).
Function	disease:The pathologies of several chronic and acute diseases of the respiratory tract involve an imbalance between the proteases of cells involved in inflammatory responses and the inhibitors of these proteases. The inflammation-mediated release of neutrophil elastase in the lungs of patients whose levels of active alpha-1-antiprotease are compromised by genetic background, cigarette smoking, air pollutants, or a combination of all three can result in severe lung damage and a decreased lifespan. The relatively small size

Nanjing BYabscience technology Co.,Ltd



of this protein, its lack of glycosylation and its stability make this protein a candidate for use as a therapeutic agent in diseases mediated by leukocyte elastase-antielastase imbalances.,function:Acid-stable proteinase inhibitor with strong affinities for trypsin, chymotrypsin, elastase, and cathepsin G. May prevent elastase-mediated damage to oral and possibly other

Background

This gene encodes a secreted inhibitor which protects epithelial tissues from serine proteases. It is found in various secretions including seminal plasma, cervical mucus, and bronchial secretions, and has affinity for trypsin, leukocyte elastase, and cathepsin G. Its inhibitory effect contributes to the immune response by protecting epithelial surfaces from attack by endogenous proteolytic enzymes. This antimicrobial protein has antibacterial, antifungal and antiviral activity. [provided by RefSeq, Nov 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night