



AMD Polyclonal Antibody

Catalog No	BYab-23359
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	PAM
Protein Name	Peptidyl-glycine alpha-amidating monooxygenase (PAM) [Includes: Peptidylglycine alpha-hydroxylating monooxygenase (PHM) (EC 1.14.17.3); Peptidyl-alpha-hydroxyglycine alpha-amidating lyase (EC 4.3.2.5)]
Immunogen	Synthesized peptide derived from human protein . at AA range: 380-460
Specificity	AMD Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	107kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle membrane ; Single-pass membrane protein . Secretory granules. .; [Isoform 1]: Membrane; Single-pass type I membrane protein.; [Isoform 2]: Membrane; Single-pass type I membrane protein.; [Isoform 3]: Secreted. Secreted from secretory granules.; [Isoform 4]: Secreted. Secreted from secretory granules.
Tissue Specificity	Epithelium,Heart,Kidney,Lung,Thalamus,Thyroid carcinoma,
Function	alternative products:Additional isoforms seem to exist,catalytic activity:Peptidylamidoglycolate = peptidyl amide + glyoxylate.,catalytic activity:Peptidylglycine + ascorbate + O(2) = peptidyl(2-hydroxyglycine) + dehydroascorbate + H(2)O.,caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,cofactor:Bind 2 copper ions per subunit; For the monooxygenase reaction.,cofactor:Zinc; for the lyase reaction.,enzyme regulation:Inhibited by

Nanjing BYabscience technology Co.,Ltd



EDTA, phenylglyoxal and diethyl pyrocarbonate.,function:Bifunctional enzyme that catalyzes 2 sequential steps in C-terminal alpha-amidation of peptides. The monooxygenase part produces an unstable peptidyl(2-hydroxyglycine) intermediate that is dismutated to glyoxylate and the corresponding desglycine peptide amide by the lyase part. C-terminal amidation of peptides such as neuropep

Background

This gene encodes a multifunctional protein. The encoded preproprotein is proteolytically processed to generate the mature enzyme. This enzyme includes two domains with distinct catalytic activities, a peptidylglycine alpha-hydroxylating monooxygenase (PHM) domain and a peptidyl-alpha-hydroxyglycine alpha-amidating lyase (PAL) domain. These catalytic domains work sequentially to catalyze the conversion of neuroendocrine peptides to active alpha-amidated products. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658