



DCAM Polyclonal Antibody

Catalog No	BYab-23943
Isotype	IgG
Reactivity	Human;Rat
Applications	WB;ELISA
Gene Name	AMD1 AMD
Protein Name	S-adenosylmethionine decarboxylase proenzyme (AdoMetDC) (SAMDC) (EC 4.1.1.50) [Cleaved into: S-adenosylmethionine decarboxylase alpha chain; S-adenosylmethionine decarboxylase beta chain]
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DCAM Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	36kD
Cell Pathway	cytosol,
Tissue Specificity	Heart,Placenta,Prostate,Thymus,Trachea,
Function	catalytic activity:S-adenosyl-L-methionine = (5-deoxy-5-adenosyl)(3-aminopropyl)-methylsulfonium salt + CO(2).,cofactor:Pyruvoyl group.,enzyme regulation:Both proenzyme processing and catalytic activity are stimulated by putrescine. Catalytic activity is inhibited by iodoacetic acid.,pathway:Amine and polyamine biosynthesis; S-adenosylmethioninamine biosynthesis; S-adenosylmethioninamine from S-adenosyl-L-methionine: step 1/1.,PTM:Is synthesized initially as an inactive proenzyme. Formation of the active enzyme involves a self-maturation process in which the active site pyruvoyl group is generated from an internal serine residue via an autocatalytic post-translational modification. Two non-identical subunits are generated from the proenzyme in this reaction, and the pyruvate is formed at the

Nanjing BYabscience technology Co.,Ltd



N-terminus of the alpha chain, which is derived from the carboxyl end of the proenzyme. The post-

Background

This gene encodes an important intermediate enzyme in polyamine biosynthesis. The polyamines spermine, spermidine, and putrescine are low-molecular-weight aliphatic amines essential for cellular proliferation and tumor promotion. Multiple alternatively spliced transcript variants have been identified. Pseudogenes of this gene are found on chromosomes 5, 6, 10, X and Y. [provided by RefSeq, Dec 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night