



HNF1A Polyclonal Antibody

Catalog No	BYab-23960
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	HNF1A TCF1
Protein Name	Hepatocyte nuclear factor 1-alpha (HNF-1-alpha) (HNF-1A) (Liver-specific transcription factor LF-B1) (LFB1) (Transcription factor 1) (TCF-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HNF1A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	69kD
Cell Pathway	Nucleus .
Tissue Specificity	Liver.
Function	disease:Defects in HNF1A are a cause of susceptibility to insulin-dependent diabetes mellitus (IDDM) [MIM:222100].,disease:Defects in HNF1A are the cause of maturity onset diabetes of the young type 3 (MODY3) [MIM:600496]; also symbolized MODY-3. MODY [MIM:606391] is a form of diabetes characterized by an autosomal dominant mode of inheritance, age of onset of 25 years or younger and a primary defect in insulin secretion. The clinical phenotype of MODY3 is characterized by severe insulin secretory defects, and by major hyperglycemia associated with microvascular complications.,disease:Defects in HNF1A may predispose to hepatic adenomas [MIM:142330]. Hepatic adenomas are benign tumors at risk of malignant transformation. Bi-allelic inactivation of HNF1A, whether sporadic or associated with MODY3, may be an early step in the developmant of some hepatocellular carcinomas.,function:Required

Nanjing BYabscience technology Co.,Ltd

**Background**

The protein encoded by this gene is a transcription factor required for the expression of several liver-specific genes. The encoded protein functions as a homodimer and binds to the inverted palindrome 5'-GTTAATNATTAAC-3'. Defects in this gene are a cause of maturity onset diabetes of the young type 3 (MODY3) and also can result in the appearance of hepatic adenomas. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Apr 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658