



ZO-2 Polyclonal Antibody

Catalog No	BYab-28986
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	TJP2
Protein Name	Tight junction protein ZO-2
Immunogen	The antiserum was produced against synthesized peptide derived from human ZO-2. AA range:1063-1112
Specificity	ZO-2 Polyclonal Antibody detects endogenous levels of ZO-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TJP2; X104; ZO2; Tight junction protein ZO-2; Tight junction protein 2; Zona occludens protein 2; Zonula occludens protein 2
Observed Band	160kD
Cell Pathway	Cell junction, adherens junction. Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell junction, tight junction . Nucleus . Also nuclear under environmental stress conditions and in migratory endothelial cells and subconfluent epithelial cell cultures. .
Tissue Specificity	This protein is found in epithelial cell junctions. Isoform A1 is abundant in the heart and brain. Detected in brain and skeletal muscle. It is present almost exclusively in normal tissues. Isoform C1 is expressed at high level in the kidney, pancreas, heart and placenta. Not detected in brain and skeletal muscle. Found in normal as well as in most neoplastic tissues.
Function	disease:Defects in TJP2 are involved in familial hypercholanemia (FHCA) [MIM:607748]. FHCA is a disorder characterized by elevated serum bile acid concentrations, itching, and fat malabsorption.,function:Plays a role in tight junctions and adherens junctions.,similarity:Belongs to the MAGUK family.,similarity:Contains 1 guanylate kinase-like domain.,similarity:Contains 1

Nanjing BYabscience technology Co.,Ltd



SH3 domain.,similarity:Contains 3 PDZ (DHR) domains.,subcellular location:Also nuclear under environmental stress conditions and in migratory endothelial cells and subconfluent epithelial cell cultures.,subunit:Homodimer, and heterodimer with ZO1. Interacts with occludin, SAFB and UBN1. Interaction with SAFB occurs in the nucleus.,tissue specificity:This protein is found in epithelial cell junctions. Isoform A1 is abundant in the heart and brain whereas isoform C1 is expressed at high level in the kidney, pancreas, heart

Background

This gene encodes a zonula occluden that is a member of the membrane-associated guanylate kinase homolog family. The encoded protein functions as a component of the tight junction barrier in epithelial and endothelial cells and is necessary for proper assembly of tight junctions. Mutations in this gene have been identified in patients with hypercholanemia, and genomic duplication of a 270 kb region including this gene causes autosomal dominant deafness-51. Alternatively spliced transcripts encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysate from K562 cells, using ZO-2 antibody.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658