



# ETAR Polyclonal Antibody

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No         | BYab-29336                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Isotype            | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Reactivity         | Human;Mouse;Monkey                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Applications       | WB;IHC;IF;ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Gene Name          | EDNRA                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Protein Name       | Endothelin-1 receptor                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human EDNRA. AA range:378-427                                                                                                                                                                                                                                                                                                                                               |
| Specificity        | ETAR Polyclonal Antibody detects endogenous levels of ETAR protein.                                                                                                                                                                                                                                                                                                                                                                             |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                         |
| Source             | Polyclonal, Rabbit,IgG                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                                                                                                                                                                                                                                                                                                                           |
| Dilution           | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.                                                                                                                                                                                                                                                                                    |
| Concentration      | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Purity             | ≥90%                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Storage Stability  | -20°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Synonyms           | EDNRA; ETA; ETRA; Endothelin-1 receptor; Endothelin A receptor; ET-A; ETA-R; hET-AR                                                                                                                                                                                                                                                                                                                                                             |
| Observed Band      | 48kD                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Cell Pathway       | Cell membrane; Multi-pass membrane protein.                                                                                                                                                                                                                                                                                                                                                                                                     |
| Tissue Specificity | Isoform 1, isoform 3 and isoform 4 are expressed in a variety of tissues, with highest levels in the aorta and cerebellum, followed by lung, atrium and cerebral cortex, lower levels in the placenta, kidney, adrenal gland, duodenum, colon, ventricle and liver but no expression in umbilical vein endothelial cells. Within the placenta, isoform 1, isoform 2, isoform 3 and isoform 4 are expressed in the villi and stem villi vessels. |
| Function           | function:Receptor for endothelin-1. Mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. The rank order of binding affinities for ET-A is: ET1 > ET2 >> ET3.,similarity:Belongs to the G-protein coupled receptor 1 family.,subunit:Interacts with HDAC7 and HTATIP.,tissue specificity:Isoform 1,                                                                          |

Nanjing BYabscience technology Co.,Ltd



isoform 3 and isoform 4 are expressed in a variety of tissues, with highest levels in the aorta and cerebellum, followed by lung, atrium and cerebral cortex, lower levels in the placenta, kidney, adrenal gland, duodenum, colon, ventricle and liver but no expression in umbilical vein endothelial cells. Within the placenta, isoform 1, isoform 2, isoform 3 and isoform 4 are expressed in the villi and stem villi vessels.,

**Background**

This gene encodes the receptor for endothelin-1, a peptide that plays a role in potent and long-lasting vasoconstriction. This receptor associates with guanine-nucleotide-binding (G) proteins, and this coupling activates a phosphatidylinositol-calcium second messenger system. Polymorphisms in this gene have been linked to migraine headache resistance. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images




Western Blot analysis of various cells using ETAR Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunofluorescence analysis of LOVO cells, using EDNRA Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from COS7 cells, using EDNRA Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human colon cancer. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).