



CD241 Polyclonal Antibody

Catalog No	BYab-30088
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	RHAG
Protein Name	Ammonium transporter Rh type A
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human RHAG. AA range:1-50
Specificity	CD241 Polyclonal Antibody detects endogenous levels of CD241 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RHAG; RH50; Ammonium transporter Rh type A; Erythrocyte membrane glycoprotein Rh50; Erythrocyte plasma membrane 50 kDa glycoprotein; Rh50A; Rhesus blood group family type A glycoprotein;Rh family type A glycoprotein; Rh type A glycoprotein; Rhesus blood group-associated ammonia channel; Rhesus blood group-associated glycoprotein; CD241
Observed Band	44kD
Cell Pathway	Membrane ; Multi-pass membrane protein.
Tissue Specificity	Erythrocytes.
Function	disease:Defects in RHAG are the cause of regulator type Rh-null hemolytic anemia (RHN) [MIM:268150]; also called Rh-deficiency syndrome. RHN is a form of chronic hemolytic anemia in which the red blood cells have a stomatocytosis and spherocytosis morphology, an increased osmotic fragility, an altered ion transport system, and abnormal membrane phospholipid organization.,function:Associated with rhesus blood group antigen expression.

Nanjing BYabscience technology Co.,Ltd



	May be part of an oligomeric complex which is likely to have a transport or channel function in the erythrocyte membrane.,similarity:Belongs to the ammonium transporter (TC 2.A.49) family. Rh subfamily.,subunit:Heterotetramer.,tissue specificity:Erythrocytes.,
Background	The protein encoded by this gene is erythrocyte-specific and is thought to be part of a membrane channel that transports ammonium and carbon dioxide across the blood cell membrane. The encoded protein appears to interact with Rh blood group antigens and Rh30 polypeptides. Defects in this gene are a cause of regulator type Rh-null hemolytic anemia (RHN), or Rh-deficiency syndrome.[provided by RefSeq, Mar 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998


监督电话: 15950492658




Products Images



Western Blot analysis of K562 cells using CD241 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100

Nanjing BYabs science technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100