



IRAK-M Polyclonal Antibody

Catalog No	BYab-30819
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	IRAK3
Protein Name	Interleukin-1 receptor-associated kinase 3
Immunogen	The antiserum was produced against synthesized peptide derived from human IRAK3. AA range:491-540
Specificity	IRAK-M Polyclonal Antibody detects endogenous levels of IRAK-M protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IRAK3; Interleukin-1 receptor-associated kinase 3; IRAK-3; IL-1 receptor-associated kinase M; IRAK-M
Observed Band	68kD
Cell Pathway	Cytoplasm . Nucleus . In dendritic cells, translocates into the nucleus upon IL33 stimulation. .
Tissue Specificity	Expressed in eosinophils, dendritic cells and/or monocytes (at protein level) (PubMed:29686383). Expressed predominantly in peripheral blood lymphocytes (PubMed:10383454).
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:Ser-293 is present instead of the conserved Asp which is expected to be an active site residue. Low level autophosphorylation activity has been reported in PubMed:10383454, while other authors describe this as an inactive kinase.,cofactor:Magnesium.,disease:Defects in IRAK3 are associated with susceptibility to asthma-related traits type 5 (ASRT5) [MIM:611064]. Asthma-related traits include clinical symptoms of asthma, such as coughing, wheezing, dyspnea, bronchial hyperresponsiveness as assessed by

Nanjing BYabscience technology Co.,Ltd



methacholine challenge test, serum IgE levels, atopy, and atopic dermatitis.,function:Inhibits dissociation of IRAK1 and IRAK4 from the Toll-like receptor signaling complex by either inhibiting the phosphorylation of IRAK1 and IRAK4 or stabilizing the receptor complex.,similarity:Belongs to the protein kinase superfamily. TK

Background

This gene encodes a member of the interleukin-1 receptor-associated kinase protein family. Members of this family are essential components of the Toll/IL-R immune signal transduction pathways. This protein is primarily expressed in monocytes and macrophages and functions as a negative regulator of Toll-like receptor signaling. Mutations in this gene are associated with a susceptibility to asthma. Alternate splicing results in multiple transcript variants. [provided by RefSeq, May 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of LOVO cells using IRAK-M Polyclonal Antibody



Immunofluorescence analysis of HeLa cells, using IRAK3 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human brain, using IRAK3 Antibody. The picture on the right is blocked with the synthesized peptide.

Nanjing BYabs science technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Western blot analysis of lysates from K562 cells, using IRAK3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVEC cells using IRAK3 antibody.