



# Neuregulin-3 Polyclonal Antibody

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No</b>         | BYab-31681                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Isotype</b>            | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Reactivity</b>         | Human;Mouse                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Applications</b>       | WB;IHC;IF;ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Gene Name</b>          | NRG3                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Protein Name</b>       | Pro-neuregulin-3 membrane-bound isoform                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Immunogen</b>          | The antiserum was produced against synthesized peptide derived from the Internal region of human NRG3. AA range:311-360                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Specificity</b>        | Neuregulin-3 Polyclonal Antibody detects endogenous levels of Neuregulin-3 protein.                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Source</b>             | Polyclonal, Rabbit,IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Dilution</b>           | WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000.. IF 1:50-200                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Concentration</b>      | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Purity</b>             | ≥90%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Storage Stability</b>  | -20°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Synonyms</b>           | NRG3; Pro-neuregulin-3, membrane-bound isoform; Pro-NRG3                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Observed Band</b>      | 80kD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Cell Pathway</b>       | [Pro-neuregulin-3, membrane-bound isoform]: Cell membrane ; Single-pass type I membrane protein . Does not seem to be active. .; [Neuregulin-3]: Secreted .; [Isoform 3]: Cell membrane; Single-pass type I membrane protein. Isoform 3 is also proteolytically released as a soluble form.                                                                                                                                                                                              |
| <b>Tissue Specificity</b> | Highly expressed in most regions of the brain with the exception of corpus callosum. Expressed at lower level in testis. Not detected in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, ovary, small intestine, colon and peripheral blood leukocytes.                                                                                                                                                                                       |
| <b>Function</b>           | developmental stage:Isoform 3 is expressed in fetal brain but not in other fetal tissues.,domain:ERBB receptor binding is elicited entirely by the EGF-like domain.,domain:The cytoplasmic domain may be involved in the regulation of trafficking and proteolytic processing. Regulation of the proteolytic processing involves initial intracellular domain dimerization.,function:Direct ligand for the ERBB4 tyrosine kinase receptor. Binding results in ligand-stimulated tyrosine |

**Nanjing BYabscience technology Co.,Ltd**



phosphorylation and activation of the receptor. Does not bind to the EGF receptor, ERBB2 or ERBB3 receptors. May be a survival factor for oligodendrocytes.,PTM:Extensive glycosylation precedes the proteolytic cleavage (By similarity). Isoform 3 is glycosylated.,PTM:Proteolytic cleavage close to the plasma membrane on the external face leads to the release of the soluble growth factor form.,similarity:Belongs to the neuregul

**Background**

This gene is a member of the neuregulin gene family. This gene family encodes ligands for the transmembrane tyrosine kinase receptors ERBB3 and ERBB4 - members of the epidermal growth factor receptor family. Ligand binding activates intracellular signaling cascades and the induction of cellular responses including proliferation, migration, differentiation, and survival or apoptosis. This gene encodes neuregulin 3 (NRG3). NRG3 has been shown to activate the tyrosine phosphorylation of its cognate receptor, ERBB4, and is thought to influence neuroblast proliferation, migration and differentiation by signalling through ERBB4. NRG3 also promotes mammary differentiation during embryogenesis. Linkage studies have implicated this gene as a susceptibility locus for schizophrenia and schizoaffective disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcri

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd


网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998


监督电话: 15950492658




## Products Images



Western Blot analysis of mouse brain, K562 cells using Neuregulin-3 Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Western Blot analysis of MOUSE-BRAIN cells using Neuregulin-3 Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:200



Western Blot analysis of mouse-heart mouse-brain mouse-lung using Neuregulin-3 Polyclonal Antibody diluted at 1:800. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



1 mouse-heart  
2 mouse-brain  
3 mouse-lung

Nanjing BYabs science technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658