



Rab11-FIP1 Polyclonal Antibody

Catalog No	BYab-31826
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	RAB11FIP1
Protein Name	Rab11 family-interacting protein 1
Immunogen	Synthesized peptide derived from Rab11-FIP1 . at AA range: 70-150
Specificity	Rab11-FIP1 Polyclonal Antibody detects endogenous levels of Rab11-FIP1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RAB11FIP1; RCP; Rab11 family-interacting protein 1; Rab11-FIP1; Rab-coupling protein
Observed Band	
Cell Pathway	Recycling endosome . Rab11A rather than Rab4A mediates localization in the endocytic recycling compartment (ERC).; [Isoform 2]: Cytoplasmic vesicle, phagosome membrane. Membrane-bound (isoform 2). Colocalizes with Rab11A at phagosomes (isoform 2).
Tissue Specificity	Isoform 2 is expressed in brain, heart, testis, lung, spleen, ovary and small intestine.
Function	function:A Rab11 effector protein involved in the endosomal recycling process. Also involved in controlling membrane trafficking along the phagocytic pathway and in phagocytosis.,similarity:Contains 1 C2 domain.,subcellular location:Membrane-bound (isoform 2). Colocalizes with Rab11A at phagosomes (isoform 2).,subcellular location:Rab11A rather than Rab4A mediates localization in the endocytic recycling compartment (ERC).,subunit:Homooligomer (isoform 2). Isoform 2 interacts with RAB4A, RAB11A, RAB11B and RAB25. According to PubMed:15280022, RAB4A binding to RAB11FIP1 is of very low affinity in vitro

Nanjing BYabscience technology Co.,Ltd



and in vivo.,tissue specificity:Isoform 2 is expressed in brain, heart, testis, lung, spleen, ovary and small intestine.,

Background

This gene encodes one of the Rab11-family interacting proteins (Rab11-FIPs), which play a role in the Rab-11 mediated recycling of vesicles. The encoded protein may be involved in endocytic sorting, trafficking of proteins including integrin subunits and epidermal growth factor receptor (EGFR), and transport between the recycling endosome and the trans-Golgi network. Alternative splicing results in multiple transcript variants. A pseudogene is described on the X chromosome. [provided by RefSeq, Dec 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658