



TWIK-2 Polyclonal Antibody

Catalog No	BYab-32114
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	KCNK6
Protein Name	Potassium channel subfamily K member 6
Immunogen	Synthesized peptide derived from TWIK-2 . at AA range: 240-320
Specificity	TWIK-2 Polyclonal Antibody detects endogenous levels of TWIK-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KCNK6; TOSS; TWIK2; Potassium channel subfamily K member 6; Inward rectifying potassium channel protein TWIK-2; TWIK-originated similarity sequence
Observed Band	33kD
Cell Pathway	Membrane ; Multi-pass membrane protein .
Tissue Specificity	Widespread expression, detected in all tissues tested except for skeletal muscle. Strongest expression in placenta, pancreas, heart, colon and spleen, lower levels detected in peripheral blood leukocytes, lung, liver, kidney and thymus. Lowest expression detected in brain.
Function	function:Exhibits outward rectification in a physiological K(+) gradient and mild inward rectification in symmetrical K(+) conditions..miscellaneous:Inhibited by internal acidification and, to a small degree, by zinc. Not inhibited by quinine, quinidine or barium..similarity:Belongs to the two pore domain potassium channel (TC 1.A.1.8) family..subunit:Homodimer .,tissue specificity:Widespread expression, detected in all tissues tested except for skeletal muscle. Strongest expression in placenta, pancreas, heart, colon and spleen, lower levels detected in peripheral blood leukocytes, lung, liver, kidney and thymus. Lowest expression detected in brain.,


Nanjing BYabscience technology Co.,Ltd



Background	This gene encodes one of the members of the superfamily of potassium channel proteins containing two pore-forming P domains. This channel protein, considered an open rectifier, is widely expressed. It is stimulated by arachidonic acid, and inhibited by internal acidification and volatile anaesthetics. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658