



PPAR delta (4G5) Mouse mAb

Catalog No	BYab-33028
Isotype	IgG1
Reactivity	Human,Rat,Mouse
Applications	WB,IHC-P
Gene Name	PPARD
Alternative Names	FAAR; NR1C2; NUC1; Peroxisome proliferative activated receptor delta
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Host	Mouse
Molecular Weight	Calculated MW: 50 kDa; Observed MW: 50 kDa
Clonality	Monoclonal Antibody
Clonality No.	4G5-2D10-2H5
Dilution	WB: 1/500-1/1000 IHC: 1/50-1/100
Immunogen	Purified recombinant protein expressed in E.coli.
Purification	Affinity Purified
Conjugation	Unconjugated
Modification	Unmodified
Form	Liquid
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Concentration	1 mg/ml
Purity	≥90%
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Background	Ligand-activated transcription factor. Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma-linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids.

Nanjing BYabscience technology Co.,Ltd



Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand.



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658