



Fibronectin (PTR1352) mouse mAb

货号	BYab-33230
同位型	IgG
应用	WB; ELISA
种属	Human;Mouse; Rat;
靶点	Fibronectin
简介	>>PI3K-Akt signaling pathway;>>Focal adhesion;>>ECM-receptor interaction;>>Regulation of actin cytoskeleton;>>AGE-RAGE signaling pathway in diabetic complications;>>Bacterial invasion of epithelial cells;>>Yersinia infection;>>Amoebiasis;>>Human papillomavirus infection;>>Pathways in cancer;>>Proteoglycans in cancer;>>Small cell lung cancer
基因名称	FN1
蛋白名称	Fibronectin
特异性	Synthesized peptide derived from human Fibronectin.AA range:2400-2477
组成	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
来源	Mouse, Monoclonal/Ig2a,kappa
稀释	WB 1:500-2000, ELISA 1:5000-20000
纯化工艺	Protein G
其他名称	FN1;FN;Fibronectin;FN;Cold-insoluble globulin;CIG
分子量	285kD
实测条带	285kD
背景	This gene encodes fibronectin, a glycoprotein present in a soluble dimeric form in plasma, and in a dimeric or multimeric form at the cell surface and in extracellular matrix. The encoded preproprotein is proteolytically processed to generate the mature protein. Fibronectin is involved in cell adhesion and migration processes including embryogenesis, wound healing, blood coagulation, host defense, and metastasis. The gene has three regions subject to alternative splicing, with the potential to produce 20 different transcript variants, at least one of which encodes an isoform that undergoes proteolytic processing. The full-length nature of some variants has not been determined. [provided by RefSeq, Jan 2016],
功能	alternative products:Additional isoforms seem to exist,developmental stage:Ugl-Y1, Ugl-Y2 and Ugl-Y3 are present in the urine from 0 to 17 years of age.,disease:Defects in FN1 are the cause of glomerulopathy with fibronectin deposits type 2 (GFND2) [MIM:601894]; also known as familial glomerular nephritis with fibronectin deposits or fibronectin glomerulopathy. GFND is a genetically heterogeneous autosomal dominant disorder characterized clinically


Nanjing BYabscience technology Co.,Ltd



	by proteinuria, microscopic hematuria, and hypertension that leads to end-stage renal failure in the second to fifth decade of life.,function:Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape. Interaction with TNR mediates inhibition of cell adhesion and neurite outgrowth
细胞定位	Secreted, extracellular space, extracellular matrix .
组织表达	Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959). Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Whole cell lysates were separated by 6% SDS-PAGE, and the membrane was blotted with anti-Fibronectin (PTR1352) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: U2OS. Predicted band size: 285 kDa. Observed band size: 285 kDa.

Nanjing BYabs science technology Co., Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658