



CCDC88C rabbit pAb

货号	BYab-33285
同位型	IgG
应用	WB
种属	Human;Mouse
靶点	CCDC88C
基因名称	CCDC88C DAPLE KIAA1509
蛋白名称	Protein Daple (Coiled-coil domain-containing protein 88C) (Dvl-associating protein with a high frequency of leucine residues) (hDaple) (Hook-related protein 2) (HkRP2)
免疫原	Synthesized peptide derived from human CCDC88C
特异性	This antibody detects endogenous levels of CCDC88C at Human, Mouse
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	223kD
功能	Required for activation of guanine nucleotide-binding proteins (G-proteins) during non-canonical Wnt signaling . Binds to ligand-activated Wnt receptor FZD7, displacing DVL1 from the FZD7 receptor and leading to inhibition of canonical Wnt signaling . Acts as a non-receptor guanine nucleotide exchange factor by also binding to guanine nucleotide-binding protein G(i) alpha (Gi-alpha) subunits, leading to their activation . Binding to Gi-alpha subunits displaces the beta and gamma subunits from the heterotrimeric G-protein complex, triggering non-canonical Wnt responses such as activation of RAC1 and PI3K-AKT signaling . Promotes apical constriction of cells via ARHGEF18 .
细胞定位	Cytoplasm . Cell junction . Enriched at apical cell junctions. .
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabs science technology Co.,Ltd



Products Images